

COMBINATORICS OF k -FAREY GRAPHS

JONAH GASTER, MIGUEL LOPEZ, EMILY REXER, ZOË RIELL, AND YANG XIAO

ABSTRACT. With an eye towards studying curve systems on low-complexity surfaces, we introduce and analyze the k -Farey graphs \mathcal{F}_k and $\mathcal{F}_{\leq k}$, two natural variants of the Farey graph \mathcal{F} in which we relax the edge condition to indicate intersection number $= k$ or $\leq k$, respectively.

The former, \mathcal{F}_k , is disconnected when $k > 1$. In fact, we find that the number of connected components is infinite if and only if k is not a prime power. Moreover, we find that each component of \mathcal{F}_k is an infinite-valence tree whenever k is even, and $\text{Aut}(\mathcal{F}_k)$ is uncountable for $k > 1$.

As for $\mathcal{F}_{\leq k}$, Agol obtained an upper bound of $1 + \min\{p : p \text{ is a prime } > k\}$ for both chromatic and clique numbers, and observed that this is an equality when k is either one or two less than a prime. We add to this list the values of k that are three less than a prime equivalent to 11 (mod 12), and we show computer-assisted computations of many values of k for which equality fails.

1. INTRODUCTION

The Farey graph \mathcal{F} is constructed as follows: the vertices

$$\left\{ \frac{p}{q} : p, q \in \mathbb{Z}, \gcd(p, q) = 1 \right\} \cup \left\{ \frac{1}{0} \right\}$$

are equipped with a determinant pairing between p/q and a/b given by

$$d_{\mathcal{F}} \left(\frac{p}{q}, \frac{a}{b} \right) = \left| \det \begin{pmatrix} p & a \\ q & b \end{pmatrix} \right|,$$

and we add edges between p/q and a/b whenever $d_{\mathcal{F}}(p/q, a/b) = 1$. The Farey graph admits a description as the ‘curve graph’ of some low-complexity surfaces, in which the vertices correspond to isotopy classes of simple closed curves, and in which edges correspond to the smallest possible intersection number on the surface—in the relevant cases, these are one or two.

Variations of the curve graph in which the edge condition is altered to take into account other intersection numbers has recently attracted attention [SS, AAC⁺, Aou, JMcM]. Motivated to study the smallest complexity nontrivial such examples, we introduce the k -Farey graphs \mathcal{F}_k and $\mathcal{F}_{\leq k}$: These graphs have the same vertex set as \mathcal{F} , but we form edges of \mathcal{F}_k (resp. $\mathcal{F}_{\leq k}$) whenever

$$d_{\mathcal{F}} \left(\frac{p}{q}, \frac{a}{b} \right) = k \quad (\text{resp. } \leq k).$$

This paper investigates combinatorial properties of \mathcal{F}_k and $\mathcal{F}_{\leq k}$. For both we exploit a natural map from the vertices of \mathcal{F} to lines in the finite plane $(\mathbb{Z}/r\mathbb{Z})^2$ (see section 3). This reduction was first used by Agol, who was motivated by exploring boundary slopes of exceptional Dehn fillings for a finite-volume cusped hyperbolic 3-manifold [Ago1, Lemma 8.2]. In our language, Agol found an upper bound for the chromatic number of $\mathcal{F}_{\leq k}$ by choosing r to be the smallest prime larger than k . For \mathcal{F}_k we find $r = k$ to be useful: when k is a prime power these lines index the components.

Date: October 21, 2018.

Key words and phrases. Farey graph, Curves on surfaces.




FIGURE 1. Pieces of \mathcal{F}_7 , colored according to the connected components.

Theorem 1. *The number of connected components of \mathcal{F}_k is given by*

$$b_0(\mathcal{F}_k) = \begin{cases} p^{\ell-1}(p+1) & \text{if } k = p^\ell, \text{ for } p \text{ a prime and } \ell > 0 \\ \infty & \text{otherwise.} \end{cases}$$

To prove connectivity of the preimages of lines when $k = p^\ell$, we identify basepoints and employ a reducing strategy to show that any point in the preimage of a line can be connected through a sequence of edges to the basepoint.

For other values of k , the proof of [Theorem 1](#) is roughly as follows. We introduce an exhaustion

$$\mathcal{F}_k^{(1)} \subset \mathcal{F}_k^{(2)} \subset \dots \subset \mathcal{F}_k$$

satisfying:

- (1) For $m > k$, if two vertices are in distinct components of $\mathcal{F}_k^{(m)}$, then they are in distinct components of $\mathcal{F}_k^{(m+1)}$.
- (2) For k not a prime power, $b_0(\mathcal{F}_k^{(m)}) < b_0(\mathcal{F}_k^{(m+1)})$ for infinitely many m .

Together, these evidently imply that $b_0(\mathcal{F}_k) = \infty$ when k is not a prime power.

The automorphism group of \mathcal{F} preserves $d_{\mathcal{F}}$, and thus $\text{Aut}(\mathcal{F}) < \text{Aut}(\mathcal{F}_k)$. Because the action of $\text{PSL}(2, \mathbb{Z})$ on \mathcal{F}_k is vertex-transitive (see [subsection 2.3](#)), it is easy to see that the connected components of \mathcal{F}_k are all isomorphic. In fact, the method of proof of [Theorem 1](#) sheds more light on the structure of each component. For instance, each component is planar (see [Proposition 4](#)), and when k is even each component is an infinite-valence tree (see [Proposition 5](#)). When k is odd, the latter statement is false; in that case \mathcal{F}_k contains triangles. Nonetheless, each vertex of \mathcal{F}_k is a cut vertex for $k > 1$ (see [Proposition 6](#)).

It is not hard to see that $\text{Aut}(\mathcal{F})$ contains $\text{PSL}(2, \mathbb{Z})$ as a finite-index subgroup. In stark contrast, a consequence of the above analysis is the following corollary:

Corollary 1. *For $k > 1$ the automorphism group $\text{Aut}(\mathcal{F}_k)$ is uncountable.*

The graph \mathcal{F}_k is the genus one version of an ‘exact k -curve graph’ associated to a closed surface of genus g , where edges indicate geometric intersection number equal to k . In these terms, [Corollary 1](#) states that the automorphism group of the exact k -curve graph in genus one is quite different than the (extended) mapping class group of the torus for $k > 1$, another low-complexity counterexample in Ivanov’s ‘Meta-Conjecture’ (see [\[Iva\]](#)). However, the automorphism group of the exact k -curve graph in higher genus remains unclear, and the methods of this paper seem unlikely to shed light on this question; in particular it seems that the exact k -curve graph is connected for $g > 1$.

In contrast, it has recently been shown that the automorphism group for the ‘at-most- k -curve graph’ is isomorphic to the mapping class group when g is sufficiently large relative to k [[AAC⁺](#)]. The analogous statement for $\mathcal{F}_{\leq k}$, namely that $\text{Aut}(\mathcal{F}_{\leq k}) \cong \text{Aut}(\mathcal{F})$, seems within reach.

We turn our attention to $\mathcal{F}_{\leq k}$. Let $p(k)$ equal the smallest prime larger than k . Agol showed the bounds $\omega(\mathcal{F}_{\leq k}) \leq \chi(\mathcal{F}_{\leq k}) \leq 1 + p(k)$ for all k (see [[Ago1](#), Lemma 8.2], cf. [[GGV](#), Theorem 8.6]), and he observed that these are equalities when k is either one or two less than a prime. As a consequence, he also observed that $\omega(\mathcal{F}_{\leq k})$ is asymptotic to k via the Prime Number Theorem [[Ago2](#)]. We add to the families of k for which Agol’s bounds are equalities:

Theorem 2. *Suppose that k is a number such that $p(k) = k + 3$, and $p(k) \equiv 11 \pmod{12}$. Then we have equality of chromatic and clique numbers $\omega(\mathcal{F}_{\leq k}) = \chi(\mathcal{F}_{\leq k}) = 1 + p(k)$.*

The known cases in which $\omega(\mathcal{F}_{\leq k}) = 1 + p(k)$ have $\omega(\mathcal{F}_{\leq k}) = k + r$, for $r = 2, 3, 4$, respectively, each for infinitely many values of k . We expect there are many other such constructions lurking in this area, and we propose the following problems about the sequence $\{\omega(\mathcal{F}_{\leq k})\}$:

Problem 1 (Exceptional k). *Characterize k for which $\omega(\mathcal{F}_{\leq k}) < 1 + p(k)$.*

Problem 2 (Big Jumps). *Let $r \geq 5$. Is it true that $\{k : \omega(\mathcal{F}_{\leq k}) \geq k + r\}$ is nonempty? infinite?*

We have collected experimental data towards both problems (see [Figure 7](#)) by analyzing values of k less than 100. Intriguingly, for [Problem 1](#) it seems that many primes that are two less than a composite are such exceptions (though $k = 37$ is curiously absent from the exceptional list). For [Problem 2](#), the data verifies nonemptiness for $r \leq 6$.

Acknowledgements. The first author thanks Ser-Wei Fu and Peter Shalen for conversations that contributed to this project. All the authors thank Ian Agol and Jeffrey Lagarias for helpful correspondence, and Tarik Aougab for his helpful ideas, enthusiasm, and support. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1439786 while the authors were in residence at the Institute for Computational and Experimental Research in Mathematics in Providence, RI, during the [Summer@ICERM 2018: Low Dimensional Topology and Geometry](#) program.

2. NOTATION AND BACKGROUND

2.1. Notation. For a graph G , we indicate the number of connected components of G by $b_0(G)$ (that is, the *zero-th Betti number* of G). We indicate the vertices of G by $V(G)$, and for $a, b \in V(G)$ we indicate the edge relation by $a \sim b$. Let $\omega(G)$ and $\chi(G)$ indicate the clique and chromatic numbers of G , respectively.

2.2. Numerators, denominators and \mathbb{Z}^2 . Vertices of \mathcal{F} are reduced fractions which evidently have numerators and denominators. It is often useful in our analysis to assume that denominators are positive; since $p/q = (-p)/(-q)$ nothing is lost in this assumption. Vertices of $x/y \in \mathcal{F}$ are also in correspondence with lines $\{(\lambda x, \lambda y) : \lambda \in \mathbb{Z}\} \subset \mathbb{Z}^2$. Each such line has two primitive elements (x, y) and $(-x, -y)$. We sometimes use (x, y) and x/y interchangeably for vertices of \mathcal{F} , taking care that this identification is 2-to-1 as remarked above.

2.3. Automorphisms. The automorphism group $\text{Aut}(\mathcal{F})$ admits a homomorphism to $\mathbb{Z}/2\mathbb{Z}$ according to whether it preserves or negates the determinant of a pair. Recall that the kernel is isomorphic to $\text{PSL}(2, \mathbb{Z})$, which acts on $V(\mathcal{F})$ via *linear fractional transformations*: the action of an element of $\text{PSL}(2, \mathbb{Z})$ on $x/y \in V(\mathcal{F})$ is given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \frac{x}{y} = \frac{ax + by}{cx + dy}.$$

It is not hard to see that $\text{PSL}(2, \mathbb{Z})$ acts transitively on $V(\mathcal{F})$.

2.4. Elementary calculations. We use the following elementary fact (see, e.g. [Hat, Lemma 2.4] for the proof).

Lemma 1. *Suppose that $a, b, n \in \mathbb{Z}$ with $\gcd(a, b) = 1$, and that (x_0, y_0) is a solution to $ax + by = n$. Then the set of integral solutions to this equation is given by $\{(x_0, y_0) + m(a, b) : m \in \mathbb{Z}\}$.*

This has an easy consequence in our setting:

Lemma 2. *Suppose that $a/b \sim x/y$ for vertices $a/b, x/y \in V(\mathcal{F}_k)$. Then the neighbors of a/b in \mathcal{F}_k all have the form $\frac{x + am}{y + bm}$ for $m \in \mathbb{Z}$.*

Proof. The points (x, y) and $(-x, -y)$ are solutions to the equations $ay - bx = k$ and $ay - bx = -k$, and the lemma follows from Lemma 1. \square

We will also make use of the following exercise:

Lemma 3. *The clique number of \mathcal{F}_k is two if k is even and three if k is odd.*

Proof. Apply an element of $\text{PSL}(2, \mathbb{Z})$ that takes a vertex in a maximum clique to $1/0$. Now the neighbors of $1/0$ have the form x/k , and any two such vertices x/k and x'/k have

$$d_{\mathcal{F}}(x/k, x'/k) = k|x - x'|.$$

First, observe that at most two such vertices can be pairwise connected, since $x < x' < x''$ implies that $|x - x''| \geq 2$. Thus $\omega(\mathcal{F}_k) \leq 3$. If k is even, then as the fractions are reduced we have $|x - x'| \geq 2$, and it is clear that $\omega(\mathcal{F}_k) = 2$. When k is odd $\{1/0, 1/k, 2/k\}$ form a clique of size three. \square

3. FROM VERTICES OF \mathcal{F} TO THE PROJECTIVE LINE OVER $\mathbb{Z}/r\mathbb{Z}$

We start by recalling the projective line over $\mathbb{Z}/r\mathbb{Z}$ (see e.g. [Her] for background). An element $(a, b) \in (\mathbb{Z}/r\mathbb{Z})^2$ is called *admissible* if $(ra, rb) = 0$ implies that $r = 0$. Denote the admissible elements by $(\mathbb{Z}/r\mathbb{Z})_{\text{adm}}^2$, and let

$$\mathcal{L}_r := (\mathbb{Z}/r\mathbb{Z})_{\text{adm}}^2 / \sim, \quad \text{where } v \sim \lambda v \text{ for } \lambda \in (\mathbb{Z}/r\mathbb{Z})^*.$$

Remark 1. When $r = p$ is prime, $\mathbb{Z}/r\mathbb{Z}$ is a field, and we have the more concise description that \mathcal{L}_r is the projectivization of the vector space $(\mathbb{Z}/p\mathbb{Z})^2$.

The vertices of \mathcal{F} are in correspondence with relatively prime pairs $(p, q) \in \mathbb{Z}^2$, modulo the equivalence relation $(p, q) \equiv (-p, -q)$. Because -1 is a unit mod r the following is well-defined:

Definition 1. The reduction $\mathbb{Z}^2 \rightarrow (\mathbb{Z}/r\mathbb{Z})^2$ induces the map

$$\phi_r : V(\mathcal{F}) \rightarrow \mathcal{L}_r .$$

We will make use of the transitive action of $\mathrm{PSL}(2, \mathbb{Z})$ on $V(\mathcal{F})$:

Lemma 4. *The action $\mathrm{PSL}(2, \mathbb{Z}) \curvearrowright V(\mathcal{F})$ descends to a transitive action on \mathcal{L}_r .*

Proof. It is easy to see that the action in the quotient is well-defined. Transitivity of the action on $V(\mathcal{F})$ immediately implies transitivity on the image $\phi_r(V(\mathcal{F})) \subset \mathcal{L}_r$. To finish, the latter is equal to \mathcal{L}_r : Choose $(x, y) \in (\mathbb{Z}/r\mathbb{Z})_{\mathrm{adm}}^2$ and let $(x, y) = \lambda(x', y')$ where $\lambda = \mathrm{gcd}(x, y)$. Note that admissibility implies that $\mathrm{gcd}(\lambda, r) = 1$, so $\lambda \in (\mathbb{Z}/r\mathbb{Z})^*$, and $(x, y) \sim (x', y')$. Now $\mathrm{gcd}(x', y') = 1$, so $\phi_r(x'/y')$ is the line through (x, y) as claimed. \square

4. CONNECTIVITY OF \mathcal{F}_k

Choosing $r = k$, we find that ϕ_k is constant on components of \mathcal{F}_k :

Lemma 5. *If $v, w \in V(\mathcal{F}_k)$ are in the same component of \mathcal{F}_k then $\phi_k(v) = \phi_k(w)$.*

Proof. Suppose that $v \sim w$, and apply an element of $\mathrm{PSL}(2, \mathbb{Z})$ that takes v to the vertex $0/1$ and w to a/b . Now since $v \sim w$ we have $|a| = k$. The reduction of this equation mod k implies that $\phi_r(w) = [(0, 1)] = \phi_r(v)$. The result follows easily. \square

It is remarkable that the preimages of ϕ_k are connected when $k = p^\ell$.

Proposition 1. *When $k = p^\ell$, the subgraph induced on $\phi_k^{-1}(L)$ is connected for all $L \in \mathcal{L}_k$.*

Proof. By [Lemma 4](#) we may assume that L is the line through $(1, 1)$. Choose as a basepoint $v = 1/1$, and let a/b be another point in $\phi_k^{-1}(L)$. Observe first that $b \neq 0$: because $\phi_r(a, b) = L = \phi_r(1, 1)$, we find that b is a unit mod r , so by [Lemma 5](#) a/b and $1/0$ are in distinct components of \mathcal{F}_k . Thus we may assume that $b > 0$. We compute a path from a/b to v in two steps: We treat first the case where $b = 1$, and when $b > 1$ we show a path exists from a/b to $c/1$ for some integer c .

If $b = 1$, then a must be congruent to $1 \pmod{p^\ell}$ since $a/b \in \phi_k^{-1}(L)$, so let $a = mp^\ell + 1$ for some integer m . One can easily check that the vertices $(mp^\ell + 1)/1$ and $((m - 1)p^\ell + 1)/1$ are incident, which gives us a path from a/b to $1/1$ by adding or subtracting multiples of p^ℓ .

Now suppose that $b > 1$. We will show that there is another vertex of $\phi^{-1}(L)$, incident to a/b , and with strictly smaller denominator. Repeating this reduction as necessary, we terminate with a path from a/b to a vertex with denominator 1 as claimed.

Let x_0/y_0 be a neighbor of a/b in \mathcal{F}_k . By [Lemma 2](#) the neighbors of a/b are given by $(x_0 + am)/(y_0 + bm)$ for $m \in \mathbb{Z}$. Observe that if b divides y_0 then the equation $d_{\mathcal{F}}(a/b, x_0/y_0) = k$ would imply b is a zero divisor in $\mathbb{Z}/k\mathbb{Z}$. However, as observed above the coordinate b is necessarily a unit. Therefore b does not divide y_0 , and the division algorithm provides two values $y = y_1$ and $y = y_2$ such that $bn + y = y_0$ and $0 < |y| < b$. Let the corresponding values of n be n_1 and n_2 .

We now have two candidates for vertices of $\phi_k^{-1}(L)$ incident to a/b with smaller denominator; it remains to check that at least one of them is a vertex in \mathcal{F}_k . In particular, we need to check that $\gcd(x_0 - an_i, y_0 - bn_i) = 1$ for some i . If $\gcd(x_0 - an_i, y_0 - bn_i) \neq 1$ then the common divisor is divisible by p , so it suffices to show that $x_0 - an_i$ and $y_0 - bn_i$ are not both divisible by p . Note that the two values n_1 and n_2 are subsequent integers. Therefore, if $(x_0 - an_i, y_0 - bn_i) \equiv (0, 0) \pmod{p}$ for $i = 1, 2$, then $(a, b) \equiv (0, 0) \pmod{p}$, a contradiction. Thus one of these values of n_i produces a vertex incident to a/b with strictly smaller denominator. \square

4.1. Calculating $b_0(\mathcal{F}_k)$ when $k = p^\ell$.

Lemma 6. *When $k = p^\ell$, we have $|\mathcal{L}_k| = p^{\ell-1}(p+1)$.*

Proof. Observe that there are $p^{2\ell} - p^{2(\ell-1)}$ admissible pairs in $(\mathbb{Z}/k\mathbb{Z})^2$. Let $\Psi : (\mathbb{Z}/k\mathbb{Z})_{\text{adm}}^2 \rightarrow \mathcal{L}_k$ be the projectivization (i.e. quotient) map. Each line $L \in \mathcal{L}_k$ has the same number of preimages in $(\mathbb{Z}/k\mathbb{Z})^2$ under Ψ . In particular, by definition

$$\Psi^{-1}(L) = \{\lambda(a, b) \mid \lambda \in (\mathbb{Z}/k\mathbb{Z})^*\},$$

for some $(a, b) \in (\mathbb{Z}/k\mathbb{Z})_{\text{adm}}^2$. Since $|(\mathbb{Z}/k\mathbb{Z})^*| = p^\ell - p^{\ell-1}$, it follows that $|\Psi^{-1}(L)| = p^\ell - p^{\ell-1}$ for any $L \in \mathcal{L}_k$, and therefore

$$|\mathcal{L}_k| = \frac{p^{2\ell} - p^{2(\ell-1)}}{p^\ell - p^{\ell-1}} = p^{\ell-1}(p+1). \quad \square$$

Together, [Lemma 5](#), [Lemma 6](#), and [Proposition 1](#) immediately imply the first half of [Theorem 1](#).

4.2. $b_0(\mathcal{F}_k) = \infty$ when k is not a prime power. We start by introducing a partial order to the vertices of \mathcal{F}_k :

Definition 2. We write $\frac{a}{b} \prec \frac{p}{q}$ when $\frac{a}{b} \sim \frac{p}{q}$ in \mathcal{F}_k and $|a| \leq |p|$ and $|b| \leq |q|$.

It is easy to see that \prec is reflexive, antisymmetric, and transitive, and so defines a partial order on $V(\mathcal{F}_k)$. When $\frac{a}{b} \prec \frac{p}{q}$ we refer to a/b as a *predecessor* of p/q and p/q as a *successor* of a/b .

Lemma 7 (Eventually Comparable). *Suppose that a/b is a vertex of \mathcal{F}_k such that $|a| + |b| > k$. Then any neighbor of a/b in \mathcal{F}_k is comparable to a/b .*

Proof. We employ the following simple observation: if m and n are of the same sign then $|m - n| = ||m| - |n||$ and $|m + n| = |m| + |n|$.

Let x/y be a neighbor of a/b in \mathcal{F}_k , i.e. $|ay - bx| = k$, with $b, y > 0$. If a and x have opposite signs then we have

$$|ay - bx| = |a|y + |x|b \geq |a| + b > k,$$

by the assumption on a and b . This contradiction implies that a and x have the same sign.

Let $s = |x| - |a|$ and $t = b - y$. If x/y and a/b are not comparable, then s and t must be non-zero and of the same sign. Thus we have

$$|ay - bx| = ||a|y - b|x|| = ||a|(b - t) - b(s + |a|)| = |t|a| + sb| = |t||a| + |s|b \geq |a| + b > k.$$

This contradiction implies that x/y and a/b are comparable. \square

In fact, it is possible to classify the possibilities for predecessors of a vertex.

Lemma 8. *The number of predecessors of a vertex are as follows:*

- (1) *If k is even, each vertex has at most one predecessor.*
- (2) *If k is odd, each vertex has at most two predecessors. Moreover, if p/q does have two predecessors a_i/b_i , then $a_1/b_1 \sim a_2/b_2$ in \mathcal{F}_k .*

Proof. If $q = 0$ then there is nothing to show; indeed in this case p/q can have no predecessors. We thus assume $q > 0$, and proceed by considering two possible cases:

- (i) the vertex $1/0$ is a predecessor, or
- (ii) the vertex $1/0$ is not a predecessor.

In case (i) we have $q = k$, and the neighbors of p/q have the form $(1 + pm)/(0 + mk)$ with $m \in \mathbb{Z}$ by Lemma 2. The inequality $|1 + pm| \leq |p|$ makes it clear that there are at most two such values of m , one of which is $m = 0$. Moreover, if there are two then we have

$$d_{\mathcal{F}} \left(\frac{1 + pm}{mk}, \frac{1}{0} \right) = |mk| = k ,$$

so that both of these predecessors are connected by an edge in \mathcal{F}_k .

For case (ii), let a_1/b_1 be a predecessor of p/q . By Lemma 2 any other incident vertex to p/q has the form $(a_1 + pm)/(b_1 + qm)$ for some integer m . For such a vertex to be a predecessor we must have $|a_1 + pm| \leq |p|$ and $|b_1 + qm| \leq q$. In this case by the triangle inequality we have

$$q \geq |b_1 + qm| \geq q|m| - b_1 \geq q(|m| - 1) ,$$

so that $|m| \leq 2$. Moreover, because b_1 and q are positive we have $m \leq 0$. This means that the three possible predecessors of p/q are $\frac{2p-a_1}{2q-b_1}$, $\frac{p-a_1}{q-b_1}$, and $\frac{a_1}{b_1}$. If the first of these is indeed a predecessor, we would have $q \geq b_1 \geq q$ and $|p| \geq |a_1| \geq |p|$, so that $a_1/b_1 = -p/q$. In this case, neither of the other two possibilities are predecessors. This narrows the possible predecessors of p/q to $\frac{p-a_1}{q-b_1}$ and $\frac{a_1}{b_1}$. Moreover, it is immediate that

$$d_{\mathcal{F}} \left(\frac{p-a_1}{q-b_1}, \frac{a_1}{b_1} \right) = k .$$

Thus we conclude that in either case (i) or (ii) the vertex p/q has at most two predecessors a_i/b_i , and if it does have two then $a_1/b_1 \sim a_2/b_2$ in \mathcal{F}_k . If k is even then Lemma 3 implies that p/q can have at most one of these as predecessors. \square

We now introduce the exhaustion

$$\mathcal{F}_k^{(1)} \subset \mathcal{F}_k^{(2)} \subset \dots \subset \mathcal{F}_k$$

referred to in the introduction.

Definition 3. The *level* of a vertex p/q is $\max\{|p|, |q|\}$. For each $m \in \mathbb{N}$, let $\mathcal{F}_k^{(m)}$ indicate the subgraph of \mathcal{F}_k induced on vertices whose level is at most m .

As a consequence of Lemma 7 and Lemma 8, we obtain:

Proposition 2 (Eventual Monotonicity). *For $m > k$, if v and w are vertices in distinct components of $\mathcal{F}_k^{(m)}$, then they are in distinct components of $\mathcal{F}_k^{(m+1)}$.*

Proof. We make two observations:

- (1) Any pair of vertices a/b and x/y at level $m + 1$ are not incident in \mathcal{F}_k , and
- (2) any vertices u_1 and u_2 that are not incident in $\mathcal{F}_k^{(m)}$ cannot both be neighbors of a vertex at level $m + 1$.

Indeed, for the first, incident vertices at level $m + 1$ are comparable by [Lemma 7](#), so suppose $a/b \prec x/y$; in either case $b = y = m + 1$ or $|a| = |x| = m + 1$ we have $|ay - bx| > k$. For the second, by [Lemma 7](#) such vertices would be comparable to the vertex at level $m + 1$ and by [Lemma 8](#) they would be incident.

Now suppose towards contradiction that $v = v_0 \sim v_1 \sim \dots \sim v_n = w$ is a shortest path in $\mathcal{F}_k^{(m+1)}$ between v and w . Evidently, there must exist a vertex v_i at level $m + 1$ in this path, since v and w are in distinct components of $\mathcal{F}_k^{(m)}$. By the first observation above, both v_{i-1} and v_{i+1} must be in $\mathcal{F}_k^{(m)}$, and by the second $v_{i-1} \sim v_{i+1}$. Thus there is a shorter path from v to w . This contradiction implies that there is no path from v to w in $\mathcal{F}_k^{(m+1)}$. \square

It follows that $m \mapsto b_0(\mathcal{F}_k^{(m)})$ is non-decreasing for $m > k$. Finally, we are able to show that this sequence increases infinitely often. Precisely,

Proposition 3. *If k is not a prime power then there exist infinitely many levels m containing vertices isolated in $\mathcal{F}_k^{(m)}$.*

Proof. Let p be a prime divisor of k such that $k = p^\ell c$ for some $\ell > 0$ and c relatively prime to p . Choose $q \equiv c \pmod{p}$ where $0 < q < p^\ell$. We claim that the vertices $(q + p^\ell n)/p^\ell$ are isolated in $\mathcal{F}_k^{(q+p^\ell n)}$ when n is such that $q + p^\ell n > k$. For such n by [Lemma 7](#) we need only check that $(q + p^\ell n)/p^\ell$ has no predecessors or successors in $\mathcal{F}_k^{(q+p^\ell n)}$.

Suppose $a/b \in \mathcal{F}_k^{(q+p^\ell n)}$ is a predecessor to $(q + p^\ell n)/p^\ell$. Then we compute

$$d_{\mathcal{F}}\left(\frac{q + p^\ell n}{p^\ell}, \frac{a}{b}\right) = qb + bp^\ell n - ap^\ell = k = p^\ell c$$

Reducing the above equation mod p^ℓ immediately yields $b \equiv 0 \pmod{p^\ell}$ since q was chosen to be coprime to p . However, a/b is a predecessor, so $b = p^\ell$. We cancel p^ℓ above and rearrange to get

$$p^\ell n - a = c - q \equiv 0 \pmod{p}$$

Thus, $a \equiv 0 \pmod{p}$ and a and b have a common factor. This contradiction implies that $(q + p^\ell n)/p^\ell$ has no predecessor.

Now suppose that a/b is a successor to $(q + p^\ell n)/p^\ell \in \mathcal{F}_k^{(q+p^\ell n)}$. Then we have $a = q + p^\ell n$ and $b > p^\ell$. Let $b = p^\ell + t$ for some positive integer t . Again we compute

$$d_{\mathcal{F}}\left(\frac{q + p^\ell n}{p^\ell}, \frac{a}{b}\right) = d_{\mathcal{F}}\left(\frac{q + p^\ell n}{p^\ell}, \frac{q + p^\ell n}{p^\ell + t}\right) = tq + tp^\ell n = t(q + p^\ell) > k$$

so these vertices cannot be incident. Thus, $(q + p^\ell n)/p^\ell$ has no successors or predecessors and is therefore isolated. \square

Remark 2. [Proposition 3](#) relies on the assumption that k is not a prime power, and the conclusion would not hold without it: for $k = p^\ell$, when m is sufficiently large there are no isolated vertices at level m .

4.3. Subgraphs of \mathcal{F}_k . We give more attention to the subgraphs of \mathcal{F}_k given by preimages under ϕ_k . Choose a line $L \in \mathcal{L}_k$.

Proposition 4. *The restriction of the natural embedding $\mathcal{F} \rightarrow \overline{\mathbb{H}}$ to the preimage $\phi_k^{-1}(L)$ is a planar embedding.*

Remark 3. See [subsection 5.1](#) for detail about the ‘natural embedding’ of \mathcal{F} ; in particular the above proposition implies that each component of \mathcal{F}_k is planar.

Proof. Suppose on the contrary that, for vertices $v, w, v', w' \in \phi_k^{-1}(L)$ with $v \sim w$ and $v' \sim w'$ in \mathcal{F}_k , we have $v < v' < w < w'$. After applying an element of $\text{PSL}(2, \mathbb{Z})$ (and switching labels if necessary) we may assume that $v = 0/1$ and $w = k/q$, and let $v' = x/y$ and $w' = a/b$.

Note that after this normalization, the line L in the image goes through $(0, 1)$, and hence for any element $c/d \in \phi_k^{-1}(L)$ we must have that k divides c . Thus the inequality $x/y < k/q < a/b$ can be simplified to

$$\frac{x'}{y} < \frac{1}{q} < \frac{a'}{b},$$

where $a = a'k$ and $x = x'k$ for $a', x' \neq 0$. This implies $a'q > b$ and $y > x'q$. Let $a'q = b + t$ and $y = x'q + s$ for positive integers s, t . Because $a/b \sim x/y$ in \mathcal{F}_k (and $x/y < a/b$ by our normalization) we have

$$k = ay - bx = (a'y - bx')k = (a'(x'q + s) - (a'q - t)x')k = (a's + x't)k > k. \quad \square$$

When k is even, we characterize the isomorphism type of the graph \mathcal{F}_k :

Proposition 5. *For k even, each component of \mathcal{F}_k is an infinite-valence tree.*

Proof. Suppose first that a component of \mathcal{F}_k contains an embedded cycle C that includes some vertex whose level is larger than k . Among the vertices of C at maximal level, let v be the vertex that is maximal with respect to \prec . Observe that the two neighbors of v in C are both comparable to it by [Lemma 7](#), and so by maximality they are both predecessors, contradicting [Lemma 8](#). Thus any embedded cycle must have its vertex levels bounded by k .

Now suppose that \mathcal{F}_k contains an embedded cycle C whose vertex levels are at most k . Choose a vertex $a/b \in C$ so that $b \neq 0$ —if C is an embedded cycle such a vertex exists. The element

$$A = \begin{pmatrix} 1 & N \\ 0 & 1 \end{pmatrix}$$

of $\text{PSL}(2, \mathbb{Z})$ acts by automorphisms of \mathcal{F}_k , taking (a, b) to $(a + bN, b)$. For N large, $A \cdot C$ is an embedded cycle of \mathcal{F}_k with a vertex whose level is larger than k , reducing to the previous case. \square

The previous lemma fails when k is odd (e.g. by [Lemma 3](#)). Nonetheless, we have:

Proposition 6. *If $k > 1$ every vertex of \mathcal{F}_k is a cut vertex.*

Proof. Again by transitivity of $\text{Aut}(\mathcal{F}) \curvearrowright V(\mathcal{F})$ we need only show this result for one vertex in \mathcal{F}_k , so choose $v = 0/1$ and let $L = \phi_k(v)$. We will show that there are no edges in $\phi_k^{-1}(L)$ between a positive vertex and a negative vertex. As $1/0 \notin \phi_k^{-1}(L)$, it follows that such vertices must be in distinct components of $\phi_k^{-1}(L) \setminus \{v\}$.

Indeed, suppose that $a/b, x/y \in \phi_k^{-1}(L)$ with $a, b, y > 0$ and $x < 0$. Immediately, we have $x \equiv a \equiv 0 \pmod{k}$, so that there are $a', x' > 0$ with $a = a'k$ and $x = -x'k$. It follows that $ay - bx = k(a'y + bx') > k$. \square

This is enough to conclude:

Proposition 7. *For $k > 1$, $\text{Aut}(\phi_k^{-1}(L))$ is uncountable.*

Proof. We claim that for each vertex v in a component $G \subset \mathcal{F}_k$, there are infinitely many components of $G \setminus \{v\}$ that are isomorphic. The claim follows as the set of permutations of \mathbb{N} is uncountable.

Again, by transitivity of $\text{Aut}(\mathcal{F}) \curvearrowright V(\mathcal{F})$ we may assume that $v = 0/1$. Note that the cyclic subgroup of $\text{PSL}(2, \mathbb{Z})$ generated by $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ fixes v . Moreover, for any $w \neq v$, the sequence $\{A^n w\}$ approaches v from the positive side and $\{A^{-n} w\}$ approaches v from the negative side.

Now choose a neighbor $w_0 \sim v$. Choosing appropriate powers of A , we find that there are $n_1, n_2 \in \mathbb{N}$ so that $A^{n_1} w_0 < 0$ and $A^{n_2} w_0 > 0$. Let $w = A^{n_1} w_0$ and $B = A^{n_2 - n_1}$ (so that $A^{n_2} w_0 = Bw$). By the proof of [Proposition 6](#), we find that w and Bw are in distinct components of $G \setminus \{v\}$, and similarly $B^n w$ and $B^{n+1} w$ are in distinct components of $G \setminus \{v\}$ for each $n \in \mathbb{Z}$. It follows that $\{B^n w\}$ picks out infinitely many components of $G \setminus \{v\}$: applying appropriate powers of B to a pair $B^{m_1} w$ and $B^{m_2} w$, we may arrange that one becomes positive and the other negative. Moreover, all such components are isomorphic, as they differ by powers of B . \square

Remark 4. It is interesting to note that there is a rich family of graph automorphisms of $\text{Aut}(\mathcal{F}_k)$ that preserves connected components of \mathcal{F}_k when $k = p^\ell$: The *level k congruence group* of $\text{PSL}(2, \mathbb{Z})$ is the kernel of $\text{PSL}(2, \mathbb{Z}) \rightarrow \text{PSL}(2, \mathbb{Z}/k\mathbb{Z})$, which induces a trivial action on the lines \mathcal{L}_k .

5. CLIQUES IN $\mathcal{F}_{\leq k}$ VIA SUBGRAPHS OF THE DUAL GRAPH TO \mathcal{F}

Agol observed that a collection of simple closed curves on the torus that pairwise intersect $\leq k$ times would have pairwise distinct images under ϕ_p [[Agol1](#), Lemma 8.2], where $p = p(k)$ is the smallest prime larger than k , from which a bound on the chromatic number is immediate. We include Agol's simple argument for completeness:

Lemma 9. *We have $\chi(\mathcal{F}_{\leq k}) \leq 1 + p(k)$.*


Proof. Suppose that $r > k$. We claim that $\phi_r : V(\mathcal{F}) \rightarrow \mathcal{L}_r$ gives a proper coloring of the vertices of $\mathcal{F}_{\leq k}$. Taking $r = p(k)$ completes the proof by [Lemma 6](#).

Suppose that $\phi_r(a/b) = \phi_r(p/q)$ for vertices $a/b, p/q \in V(\mathcal{F})$. This implies that $(p, q) \equiv \lambda(a, b)$ for $\lambda \in (\mathbb{Z}/r\mathbb{Z})^*$, and $aq \equiv bp \pmod{r}$ follows readily. In particular, we have $d_{\mathcal{F}}(a/b, p/q) = nr$ for a non-negative integer n . But $d_{\mathcal{F}}(v, w)$ is non-zero when $v \neq w$. Thus $d_{\mathcal{F}}(a/b, p/q) \geq r > k$ and we conclude $a/b \not\sim p/q$. \square

Remark 5. The proof above actually implies that $\chi(\mathcal{F}_{\leq k})$ is at most $\min\{|\mathcal{L}_r| : r > k\}$. It is unclear where there exists k for which this is smaller than $1 + p(k)$.

Since $\omega(G) \leq \chi(G)$ for all graphs G , it remains to find lower bounds for $\omega(\mathcal{F}_{\leq k})$, i.e. constructions of large cliques in $\mathcal{F}_{\leq k}$. Evidently, any collection S of vertices of \mathcal{F} will determine a clique in $\mathcal{F}_{\leq k}$ when k is large enough—in particular, when k is at least the maximum determinant pairing among elements of S . The challenge in finding good lower bounds for $\omega(\mathcal{F}_{\leq k})$ is to make sure this ‘large’ is small relative to the number of vertices of S .

We now describe how to organize our search for families of vertices according to their relative positions in \mathcal{F} . We start by recalling the relationship between the Farey graph \mathcal{F} , the hyperbolic plane \mathbb{H} , and continued fraction expansions.


 FIGURE 2. The dual graph \mathcal{G} to \mathcal{F} , and a horocycle $H_{0/1}$.


5.1. The natural planar embedding of the Farey graph. The index-two subgroup $\mathrm{PSL}(2, \mathbb{Z})$ of $\mathrm{Aut}(\mathcal{F})$ is isomorphic to a subgroup of the isometries of the hyperbolic plane \mathbb{H} . This is no accident: The Farey graph may be embedded in a compactification of the hyperbolic plane $\overline{\mathbb{H}} = \mathbb{H} \cup \partial_\infty \mathbb{H}$, in which $V(\mathcal{F})$ is in natural correspondence with $\mathbb{Q} \cup \{\infty\} \subset \mathbb{R} \cup \{\infty\} \cong \partial_\infty \mathbb{H}$ and edges are realized as hyperbolic geodesics between points of $\partial_\infty \mathbb{H}$. We refer to this as the *natural embedding* of \mathcal{F} in the plane. One manifestation of the usefulness of this point of view is the following:

The embedding $\mathcal{F} \subset \overline{\mathbb{H}}$ determines a tiling of \mathbb{H} by ideal hyperbolic triangles, and the dual graph \mathcal{G} to this tiling is a trivalent tree in \mathbb{H} . The complementary regions of \mathcal{G} are approximate horoballs—the boundary of the complementary regions is piecewise geodesic rather than horocyclic—and each region accumulates onto a single point in $V(\mathcal{F}) \subset \partial_\infty \mathbb{H}$, the ‘center’ of the horoball. We indicate the union of the edges of \mathcal{G} incident to the (approximate) horoball centered at p/q by $H_{p/q} \subset \mathcal{G}$.

Note that when $H_{p/q}$ and $H_{a/b}$ intersect we must have $p/q \sim a/b$, i.e. these vertices have determinant pairing 1. More generally, disjoint sets $H_{p/q}$ and $H_{a/b}$ are connected by a unique geodesic path in the tree \mathcal{G} , and the planar embedding $\mathcal{G} \subset \mathbb{H}$ allows us to interpret this geodesic as a sequence of paths contained in horocycles, interspersed by left- and right-turns in \mathcal{G} . We refer to the sequence of positive integers so obtained as a ‘left-right sequence’ associated to the geodesic in \mathcal{G} . In fact we have:

Theorem 3. *Suppose that (l_1, \dots, l_m) is a left-right sequence for a geodesic from $H_{p/q}$ to $H_{a/b}$, for vertices $p/q, a/b \in V(\mathcal{F})$. Then the determinant $d_{\mathcal{F}}(p/q, a/b)$ is given by the numerator of the continued fraction expansion*

$$l_1 + \frac{1}{l_2 + \frac{1}{\dots + \frac{1}{l_m + 1}}}$$

FIGURE 3. The computation of $d_{\mathcal{F}}(1/3, -2/1) = 7$.

The action $\mathrm{PSL}(2, \mathbb{Z}) \curvearrowright V(\mathcal{F})$ is transitive, $d_{\mathcal{F}}$ -preserving, and induces an automorphism that preserves left-right sequences of geodesics in \mathcal{G} . Thus for the proof of this theorem it suffices to check the claim when $a/b = 1/0$. In this case, $d_{\mathcal{F}}(1/0, p/q) = q$, and the claim follows from [Hat, Thm. 2.1, p. 31]. We encourage readers to consult [Hat] for a beautiful detailed exposition.

Example 1. The geodesic from $H_{-2/1}$ to $H_{1/3}$ is described by both left-right sequences $\{2, 2\}$ and $\{1, 1, 2\}$ (see Figure 3); as well the geodesic from $H_{1/3}$ to $H_{-2/1}$ is given by $\{3, 1\}$ and $\{1, 2, 1\}$. Note that we have

$$\begin{aligned} \boxed{2} + \frac{1}{\boxed{2}+1} &= \frac{\boxed{7}}{3}, & \boxed{1} + \frac{1}{\boxed{1} + \frac{1}{\boxed{2}+1}} &= \frac{\boxed{7}}{4}, \\ \boxed{3} + \frac{1}{\boxed{1}+1} &= \frac{\boxed{7}}{2}, & \text{and } \boxed{1} + \frac{1}{\boxed{2} + \frac{1}{\boxed{1}+1}} &= \frac{\boxed{7}}{5}, \end{aligned}$$

and $d_{\mathcal{F}}(-2/1, 1/3) = 7$.

The following observation is immediate from the calculation of Theorem 3.

Lemma 10. *If $\{a_1, \dots, a_m\}$ and $\{b_1, \dots, b_m\}$ are two left-right sequences with $a_i \leq b_i$ for each i , then the intersection number determined by $\{a_i\}$ is at most that determined by $\{b_i\}$.*


Given a finite subgraph $K \subset \mathcal{G}$, consider the associated subset $V_K \subset V(\mathcal{F})$ determined by

$$V_K = \left\{ \frac{p}{q} : H_{p/q} \cap K \text{ is neither empty nor a singleton} \right\},$$

and we let

$$I(K) = \max \left\{ d_{\mathcal{F}} \left(\frac{p}{q}, \frac{a}{b} \right) : \frac{p}{q}, \frac{a}{b} \in V_K \right\}.$$

See Figure 4 for an example.


 FIGURE 4. A subgraph K of \mathcal{G} with $I(K) = 5$ and $|V_K| = 6$.

We now describe several explicit constructions of finite subgraphs of \mathcal{G} . See Figure 5 for visual presentations.

Definition 4. Let P_n indicate an embedded path of length n inside a horocycle of \mathcal{G} .

- (1) Let R_n be the union of P_n with the $n - 1$ edges of \mathcal{G} incident to non-endpoint vertices of P_n .
- (2) There is a vertex of R_{2n} that is determined by the property that it is the unique fixed endpoint of an involution of R_{2n} . Let S_n be the union of R_{2n} with the two edges of \mathcal{G} incident to this endpoint.

Lemma 11. We have $I(R_n) = n - 1$, $|V_{R_n}| = n + 1$, $I(S_n) = 2n - 1$, and $|V_{S_n}| = 2n + 2$.

Proof. The left-right sequences between non-neighboring horoballs incident to R_n are given by $\{\{i\} : 1 \leq i \leq n - 2\}$. The extreme scenario is evidently $\{n - 2\}$, so $I(R_n) = n - 1$ by Lemma 10.

For S_n , a set of left-right sequences for pairs of non-neighboring horoballs is given by $\{\{i\} : 1 \leq i \leq 2n - 2\} \cup \{\{j, 1\} : 1 \leq j \leq n - 1\}$. It is not hard to see that both extreme cases give $I(S_n) = 2n - 1$. \square

The third construction is the most involved. Start with P_{7n-1} , and number the non-endpoint vertices of P_{7n-1} in order $1, \dots, 7n - 2$. Now decompose these vertices into the sets

$$\begin{aligned} J_1 &= \{1, 2, \dots, 2n - 1\} \cup \{5n, 5n + 1, \dots, 7n - 2\}, \\ J_2 &= \{2n, 2n + 1, \dots, 3n - 1\} \cup \{4n, 4n + 1, \dots, 5n - 1\}, \text{ and} \\ J_3 &= \{3n, 3n + 1, \dots, 4n - 1\}. \end{aligned}$$


For each vertex $v \in J_1$, add the unique edge of $\mathcal{G} \setminus P_{7n-1}$ incident to v . For each vertex $v \in J_2$, add the three edges of $\mathcal{G} \setminus P_{7n-1}$ that are within distance one of v . Finally, for each $v \in J_3$, add the seven edges of $\mathcal{G} \setminus P_{7n-1}$ that are within distance two of v . We refer to the result as T_n .

Lemma 12. We have $I(T_n) = 12n - 4$, and $|V_{T_n}| = 12n$.

Proof. By Lemma 10, it is enough to check the extreme left-right sequences for pairs of non-neighboring horoballs incident to T_n . It is not hard to see that such extremes are given by the left-right sequences

$$\begin{aligned} &\{7n - 3\}, \{5n - 2, 1\}, \{4n - 2, 2\}, \{4n - 2, 1, 1\}, \\ &\{2, 3n - 2, 1\}, \{2, 2n - 2, 2\}, \{2, 2n - 2, 1, 1\}, \\ &\{2, 1, n - 2, 2\}, \{2, 1, n - 2, 1, 1\}, \{3, n - 2, 2\}, \{3, n - 2, 1, 1\}. \end{aligned}$$

These sequences determine continued fraction expansions with numerators $7n - 2, 10n - 3, 12n - 5, 12n - 4, 12n - 4, 12n - 7, 12n - 5, 9n - 9, 9n - 6, 9n - 12$, and $9n - 9$, respectively, so that $I(T_n) = 12n - 4$, as claimed. \square

FIGURE 5. Subgraphs of \mathcal{F} with small maximum intersection.

Proposition 8. *We have:*

- (1) For all k , $\omega(\mathcal{F}_{\leq k}) \geq k + 2$.
- (2) For k odd, $\omega(\mathcal{F}_{\leq k}) \geq k + 3$.
- (3) For $k \equiv 8 \pmod{12}$, $\omega(\mathcal{F}_{\leq k}) \geq k + 4$.

Proof. These are immediate consequences of [Lemma 11](#) and [Lemma 12](#): For (1), we have $I(R_{k+1}) = k$ and $|V_{R_{k+1}}| = k + 2$; for (2) taking $k = 2n - 1$ we have $I(S_n) = k$ and $|V_{S_n}| = k + 3$; finally for (3) if $k = 12n - 4$ then $I(T_n) = k$ and $|V_{T_n}| = k + 4$. \square

[Theorem 2](#) follows immediately.

5.2. Computational data for $\omega(\mathcal{F}_{\leq k})$. There are many existing computational tools for computing graph invariants, including of course clique number. Using Mathematica to search through large finite subgraphs of $\mathcal{F}_{\leq k}$, we obtained the data presented in [Figure 7](#). The values of k presented are exactly those, less than 100, for which experimental data suggests that $\omega(\mathcal{F}_{\leq k}) < 1 + p(k)$. We make several comments about the data:

- (1) The list appears to include many primes that are two less than a composite. Indeed, among $k \leq 100$ the only exception is $k = 37$.
- (2) While the values of the third column of [Figure 7](#) might be reduced in light of [Remark 5](#), this observation will not resolve the gaps between the second and third columns. Indeed, for $k = 7$, we have $\min\{|\mathcal{L}_r| : r > 7\} = 12 = 1 + p(7)$, while the largest clique found experimentally in $\mathcal{F}_{\leq 7}$ has size 10.
- (3) It would be interesting to investigate the chromatic numbers $\chi(\mathcal{F}_{\leq k})$ for the values of k listed. However, initial attempts using Mathematica's software were insufficient to narrow the given bounds, even for $k = 7$.

- (4) For values of k less than 100 that are missing in the first column, experimental evidence produced cliques of size $1 + p(k)$. For instance, we have $\omega(\mathcal{F}_{\leq 24}) = 30$ (see Figure 6) and $\omega(\mathcal{F}_{\leq 48}) = 54$. These represent the largest gaps observed (namely, size six) between k and $\omega(\mathcal{F}_{\leq k})$ (cf. Problem 2).

We consider these phenomena to be worthy of further investigation.




FIGURE 6. A clique of size $30 = 1 + p(24)$ in $\mathcal{F}_{\leq 24}$.

k	Largest clique	Smallest coloring	k	Largest clique	Smallest coloring
7	10	12	67	70	72
13	16	18	73	76	80
19	23	24	74	78	80
23	27	30	75	78	80
31	34	38	79	82	84
32	36	38	83	87	90
33	37	38	84	89	90
43	46	48	85	89	90
47	51	54	89	93	98
53	57	60	90	94	98
54	59	60	91	94	98
61	65	68	92	96	98
62	67	68	93	96	98
63	67	68	97	100	102

FIGURE 7. Exceptional values of k that are less than 100 for which we are unable to decide whether $\omega(\mathcal{F}_{\leq k}) = \chi(\mathcal{F}_{\leq k})$.

REFERENCES

- [Ago1] Ian Agol. Bounds on exceptional Dehn filling. *Geom. Topol.* **4**(2000), 431–449.
- [Ago2] Ian Agol. Intersection of curves on the torus. Unpublished note.
- [AAC⁺] Shuchi Agrawal, Tarik Aougab, Yassin Chandran, Marissa Loving, John Oakley, Roberta Shapiro, and Yang Xiao. Automorphisms of k -Curve Graphs. In preparation.
- [Aou] Tarik Aougab. Local geometry of the k -curve graph. *Trans. Amer. Math. Soc.* **370**(2018), 2657–2678.
- [GGV] Jonah Gaster, Joshua Evan Greene, and Nicholas G. Vlamis. Coloring curves on surfaces. *Forum of Mathematics, Sigma* **6**(2018).
- [Hat] Allen Hatcher. Topology of numbers. *Unpublished manuscript, in preparation* (2017).
- [Her] Armin Herzer. Chain geometries. In *Handbook of Incidence Geometry*, pages 781–842. Elsevier, 1995.
- [Iva] Nikolai V. Ivanov. Automorphism of complexes of curves and of Teichmüller spaces. *Internat. Math. Res. Notices* **14**(1997), 651–666.
- [JMcM] M. Juvan, A. Malnič, and B. Mohar. Systems of curves on surfaces. *J. Combin. Theory Ser. B* **68**(1996), 7–22.
- [SS] Paul Schmutz Schaller. Mapping class groups of hyperbolic surfaces and automorphism groups of graphs. *Compositio Math.* **122**(2000), 243–260.

DEPARTMENT OF MATHEMATICS, MCGILL UNIVERSITY, BURNSIDE HALL, 805 SHERBROOKE STREET WEST,
MONTREAL, QUEBEC H3A 0B9

Email address: `jbgaster@gmail.com`,

MATH DEPARTMENT, BOSTON UNIVERSITY, 111 CUMMINGTON MALL, BOSTON, MA 02215

Email address: `mlopez3@bc.edu`,

MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, MAIL STOP: 1131-002-1AC, ATLANTA, GA 30322

Email address: `emily.rexer@gmail.com`,

DEPARTMENT OF MATHEMATICS AND STATISTICS, SMITH COLLEGE, BURTON HALL 115, NORTHAMPTON, MA
01063

Email address: `zoe.riell@gmail.com`,

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, BOX 1917, 151 THAYER STREET, PROVIDENCE, RI 02912

Email address: `yang_xiao@brown.edu`,